



PAPER 5505				
No	Working	Answer	Mark	Notes
1	(a)(i) (ii) (b)	2.21 0.013 663	2  2	B1 for 2.21 B1 for 0.013 M1 for $3 \times 13 \times 17$ oe A1 for 663
2		Tick boxes 1,3 & 6	3	B1 for $\frac{\pi abc}{2d}$ ; B1 for $2a^2$ ; B1 for $2(c^2 + d^2)$ (-B1 for each additional expression ticked (>3) to a minimum of 0)
3	(a)	$200 \times 0.2$	2	M1 for $200 \times 0.2$ or $\frac{40}{200}$ seen
	(b)	$0.2 + 0.4$	2	A1 for 40 M1 for $0.2 + 0.4$ A1 for 0.6
4	(a)	eg $2 \overline{)108}$ $2 \overline{)54}$ $3 \overline{)27}$ $3 \overline{)9}$ 3	3	M2 for full systematic method of at least 4 divisions by prime numbers oe factor trees; condone 1 calculation error (M1 for 108 written as a correct product with only one non-prime or equivalent division or a full process with 2 calculation errors.) A1 for $2^2 \times 3^3$ (accept $2 \times 2 \times 3 \times 3 \times 3$ )
	(b)	12	1	B1 for 12
5		Perpendicular from P to intersecting arcs (within tramlines); perpendicular at least 2 cm long	2	M1 relevant pair of arcs crossing within tramlines A1 SC M1A0 for full construction of a line perpendicular to AB not through P
6		$15 \times 10$	3	M1 for $15 \times 10$ A1 for 150 B1 for $\text{cm}^3$

PAPER 5505				
No	Working	Answer	Mark	Notes
7	(a)	$k^3$ .	1	B1 for $k^3$ .
	(b)(i)	$4x + 20 + 3x - 21$	4	M1 for <b>three</b> of 4 terms $4x + 20 + 3x - 21$ (or better) A1 for $7x - 1$
	(ii)	$x^2 + 3yx + 2xy + 6y^2$		M1 for <b>three</b> of 4 terms $x^2 + 3yx + 2xy + 6y^2$ A1 for $x^2 + 5xy + 6y^2$
	(c)	$(p + q)(p + q + 5)$	1	B1 for $(p + q)(p + q + 5)$
	(d)	$m^8$	1	B1 for $m^8$ .
	(e)	$6r^3t^6$	2	B2 for $6r^3t^6$ (B1 for $\dots r^3t^6$ or for $6\dots t^6$ )
8	(i)	Least length = 100.5	2	B1 for 100.5
	(ii)	Greatest length = 101.5		B1 for 101.5; ACCEPT 101.499 or better
9		24	5	B1 for $6\frac{2}{5} = \frac{32}{5}$ oe or $3\frac{1}{5} = \frac{16}{5}$ oe or $\frac{30}{8} + \frac{2}{8}$ oe (or implied by area of triangle=2) M1 for $\frac{1}{2} \times \frac{5}{8} \times 6\frac{2}{5}$ oe M1 for (area of square) = $18 \times$ product of two lengths A1 = $\sqrt{18 \times 2}$ A1 for 24

**PAPER 5505**

No	Working	Answer	Mark	Notes
10 (a)	$6y = 15 - 5x$	$y = \frac{15 - 5x}{6}$	2	M1 for either $6y = 15 - 5x$ or for $\frac{5x}{6} + y = \frac{15}{6}$ or $-6y = 5x - 15$ or a correct ft on sign error to $y =$ A1 for $y = \frac{15 - 5x}{6}$ oe
(b)	$6k + 5(-21) = 15$	20	2	M1 for subst. of $x = -21$ (or $x = 21$ ) into given eq <sup>n</sup> or answer to (a) A1 for $k = 20$
(c)(i)		Region R indicated	3	B2 correct region shaded (accept unshaded if R clear) (B1 shaded (R) region satisfies 3 of the 4 given inequalities with same boundaries)
(ii)		(1,1)		B1 for (1,1)
11 (a)		$y = 2x + 6$	2	B2 for $y = 2x + 6$ (B1 for $y = 2x + k$ , $k \neq 1$ or for $y = mx + 6$ , $m \neq 0$ or for $2x + 6$ )
(b)	Grad of $AB = 2$ ; Grad of $BC = \frac{-1}{"2"}$	$y = -\frac{1}{2}x + 6$	2	M1 for Grad of $BC = \frac{-1}{"2"}, -\frac{1}{2}$ or grad of $BC = -\frac{1}{\text{grad of } AB}$ A1 ft for $y = -\frac{1}{2}x + 6$ oe ft on wrong coeff. of $x$ in (a)
(c)	Eg A rectangle is always a cyclic quadrilateral because the opposite angles of a rectangle always add up to $180^\circ$ ( $90+90$ )		1	B1 for valid explanation. (eg lines from the pt of int. of diagonals of rect to all 4 vertices are equal (radii))
12	Vertices of new triangle at $(-1, -4)$ $(-4, 2)$ and $(2, 2)$	Correct triangle drawn	3	B3 cao (B2 for 3 correct vertices no triangle or triangle with 2 correct vertices (B1 for triangle with two of six co-ordinates correct from using P as centre or any isosceles triangle with base 6 and height 6)

**PAPER 5505**

No	Working	Answer	Mark	Notes
13 (a) (b) (c)	 <p data-bbox="347 414 1041 491">Median(B) &gt; Median(G); on average boys take longer IQR(B) &gt; IQR(G); times for boys have a greater spread</p>	32	1 3 2	B1 for 32 (accept 31.5 to 33.5 inclusive) B1 for ends of whiskers at 9 and 57 (with a box) B1 for ends of box at 16 and 45/46 ( $\pm 0.5$ ) B1 for median marked at "32" or complete box and whisker diagram drawn with a median inside the box B1 eg for comparison of medians (ft on diagrams) B1 eg for comparison of (interquartile) ranges (ft on diagram)
14	Reading top to bottom frequencies are 20; 18; 45; 52		2	B2 all correct (B1 for one frequency correct)
15 (a) (b)	$1600 = pq^0$ ; $400 = pq^2$ $q^2 = 0.25$ $V = pq^{-2}$	$p = 1600$ $q = 0.5$  $6400$	3 2	M1 for either $400 = pq^2$ or $1600 = pq^0$ B1 for $p = 1600$ A1 for $q = 0.5$ M1 for recognition that $t = -2$ A1 for 6 400

PAPER 5505				
No	Working	Answer	Mark	Notes
16	(a) (I) $AB = AC$ (triangle $ABC$ is isosceles) (II) $PB = PC$ tangents (from a point to a circle are) equal (III) $AP = AP$ (common side) so the 2 triangles are congruent ,SSS.		3	B3 for I, II, III with congruency reason (B2 for any two of I, II or III) (B1 for any one of the I, II or III)
	(b) $BPC = 20^\circ$ $PBC$ (or $PCB$ ) = $90 - 1/2$ "20" (= $80^\circ$ ) $BAC = PBC =$ "80"	$50^\circ$	4	B4 for $50^\circ$ (B3 for $BAC = 80^\circ$ ) (B2 for $PBC = 80^\circ$ or $PCB = 80^\circ$ ) (B1 for $APC = 10^\circ$ or $BPC = 20^\circ$ or a middle angle = $90^\circ$ ) SC if clear numerical slip seen eg " $PBC = 180 - 90 - 10 = 70$ " then goes on to get correct ft angle $ABC = 55$ deduct 1 from total so this cand would get $B4 - 1 = B3$
17	(a) $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ $= \overrightarrow{OA} + \frac{2}{3} (6\mathbf{c} - 6\mathbf{a})$ $= 6\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}$	$2\mathbf{a} + 4\mathbf{c}$	3	M1 for $\overrightarrow{OP} = \overrightarrow{OA} + \overrightarrow{AP}$ or any correct vector journey involving $\overrightarrow{OP}$ M1 for $\overrightarrow{AP} = \frac{2}{3} (6\mathbf{c} - 6\mathbf{a})$ oe or $\overrightarrow{CP} = \frac{1}{3} (-6\mathbf{c} + 6\mathbf{a})$ oe or reverse vectors A1 for $2\mathbf{a} + 4\mathbf{c}$ oe (accept unsimplified)
	(b) Eg $\overrightarrow{OM} = \overrightarrow{OC} + \overrightarrow{CM} = 6\mathbf{c} + 3\mathbf{a}$ $\overrightarrow{OM} = 1.5\overrightarrow{OP}$	$\overrightarrow{OM} = 1.5\overrightarrow{OP}$ so $OPM$ is a straight line	2	B1 for $\overrightarrow{OM} = 6\mathbf{c} + \frac{1}{2}(6\mathbf{a})$ or $\overrightarrow{PM} = 2\mathbf{c} + \mathbf{a}$ unsimplified or reverse vectors B1 for a fully correct proof.

PAPER 5505				
No	Working	Answer	Mark	Notes
18 (a)		4	1	B1 for 4 condone $\pm 4$
(b)		2	1	B1 for 2 condone $\pm 2$
(c)	$\sqrt{160} = 4\sqrt{10}$ ; $\left[ \frac{\sqrt{8}(\sqrt{5} + \sqrt{20}) - \sqrt{2} \times \sqrt{5}}{\sqrt{8}(\sqrt{5} + \sqrt{20})} \right] \times 100$ $\left[ \frac{6\sqrt{10} - \sqrt{10}}{6\sqrt{10}} \right] \times 100$	$\frac{500}{6}$	4	B1 for <b>either</b> $\sqrt{160} = 4\sqrt{10}$ or $\sqrt{8} = 2\sqrt{2}$ or $\sqrt{20} = 2\sqrt{5}$ M1 for $\left[ \frac{\sqrt{8}(\sqrt{5} + \sqrt{20}) - \sqrt{2} \times \sqrt{5}}{\sqrt{8}(\sqrt{5} + \sqrt{20})} \right]$ oe ( $\times 100$ ) B1 for either $6\sqrt{10} - \sqrt{10}$ or $6\sqrt{10}$ A1 for $\frac{500}{6}$ (accept 83.3 if no obvious earlier error)

PAPER 5505				
No	Working	Answer	Mark	Notes
19 (a)(i)		$(2x - 7)(x - 14)$	3	M1 $x^2$ term and constant term ( $\pm 98$ ) obtained or $2x(x-14) - 7(x-14)$ or $x(2x-7) - 14(2x-7)$ A1 for $(2x - 7)(x - 14)$ B1ft ft (i) provided of form $(2x \pm a)(x \pm b)$
(ii)		$x = \frac{7}{2}; x = 14$		
(b)(i)		$\frac{7}{n+7}$	3	B1 for $\frac{7}{n+7}$ oe
(ii)	$\frac{7}{n+7} = \frac{2}{5} \Rightarrow 2(n+7) = 5 \times 7$ $2n = 21$	$n=10.5$ is not possible since $n$ has to be an integer		M1 for $2(n+7) = 5 \times 7$ or $n+7 = 5 \times 3.5$ (can be implied) ft (b)(i) fractional in terms of $n$ and $< 1$ A1 ft for $n=10.5$ not possible (since $n$ not integer) oe
(c)	$2 \times \left(\frac{n}{n+7}\right) \times \left(\frac{7}{n+7}\right) = \frac{4}{9}$ $14n \times 9 = 4(n+7)^2$ $14n \times 9 = 4(n^2 + 14n + 49)$ $4n^2 + 56n + 196 - 126n = 0$	$2n^2 - 35n + 98 = 0$	5	M1 for $\left(\frac{n}{n+7}\right) \times \left(\frac{7}{n+7}\right)$ seen M1 for $2 \times \left(\frac{n}{n+7}\right) \times \left(\frac{7}{n+7}\right)$ oe ( $= \frac{4}{9}$ ) M1(dep on 1 <sup>st</sup> M) elimination of fractions within an equation B1 3 terms correct in expansion of $(n+7)^2 = n^2 + 7n + 7n + 49$ A1 full valid completion to printed answer

PAPER 5505				
No	Working	Answer	Mark	Notes
(d)	$\frac{7}{n+7} \times \frac{7}{n+7} = \frac{7}{21} \times \frac{7}{21} =$	$\frac{1}{9}$	2	M1 for $\frac{7}{n+7} \times \frac{7}{n+7}$ or better or ft [answer (b)(i)] <sup>2</sup> or $1 - \frac{4}{9} - \left(\frac{n}{n+7}\right)^2$ A1 for $\frac{1}{9}$ oe cao
20	(a)(i) (ii)  (b)  Stretch parallel to y-axis scale factor 3 Stretch parallel to x-axis scale factor $\frac{1}{2}$	$y = 1 + \sin x$ $y = 2 \sin x$	2  3	B1 for $y = 1 + \sin x$ B1 for $y = 2 \sin x$ SC both (i) $f(x) + 1$ , (ii) $2f(x)$ B1 M1 for 'stretch' A1 for Stretch parallel to y-axis scale factor 3 oe A1 for Stretch parallel to x-axis scale factor $\frac{1}{2}$ oe  SC if M0 award BI for "sf 3 vertically" <b>and</b> "sf $\frac{1}{2}$ horizon."