


June 2010

1380/3H				
Question	Working	Answer	Mark	Notes
1		$8x + 6y$	2	B2 for $8x + 6y$ or $6y + 8x$ or $2(4x + 3y)$ or $2(3y + 4x)$ ; accept $x8$ or $y6$ etc. [B1 for $8x$ or $6y$ , accept $x8$ or $y6$ ]
2		$\begin{array}{r l} 4 & 3\ 5\ 7\ 7 \\ 5 & 0\ 3\ 3\ 5\ 6\ 7\ 8\ 8\ 8 \\ 6 & 1\ 2\ 2 \end{array}$ <p>Key Eg. <math>4\   \ 3</math> means 43g</p>	3	B2 for a fully correct diagram. Accept a stem of 40, 50, 60 (the order of the numbers in the stem may be reversed) (B1 for ordered or unordered leaves, with just one error or omission) B1 for a correct key (units may be omitted)
3	(i) $180 - 110 = 70$ $180 - 2 \times 70$  (ii)	$40$  Reasons	4	M1 for $180 - 110$ or 70 seen M1 for $180 - 2 \times "70"$ or $110 - "70"$ A1 cao B1 for two out of three of: angles on a line add to $180^\circ$ ; isosceles triangle (accept 2 sides equal or 2 angles equal) ; sum of the angles in a triangle is equal to $180^\circ$ <b>OR</b> for two out of three of: angles on a line add to $180^\circ$ ; isosceles triangle (accept 2 sides equal or 2 angles equal) ; exterior angle of a triangle is equal to the sum of the interior opposite angles

Question		Working	Answer	Mark	Notes
4	(a)		10 10	1	B1 for 10 10 or equivalent
	(b)		13 – 14	1	B1 for answer in range 13-14 inclusive
	(c)		Line from (11 10, 40) to (11 50, 0)	1	B1 for a line drawn from (11 10, 40) to (11 50, 0) Accept a tolerance of $\pm 2$ mm
5			Triangle at (1,-2), (-1,-2), (1,-5)	2	B2 for triangle at (1,-2), (-1,-2), (1,-5) (B1 for rotation of $180^\circ$ about the wrong centre or for a rotation of $90^\circ$ , centre (1,0) clockwise or anticlockwise)
6			Enlargement scale factor 2 centre (1,0)	3	B1 for enlargement B1 for scale factor 2 oe (eg 'x2', ' by 2', ' of 2') B1 for (1,0) [condone omission of brackets and/or the word 'centre'; do not accept a vector]  Note: A combination of transformations gets NO marks

Question	Working	Answer	Mark	Notes
7	$40 \div (2 + 3) = 8$ $8 \times 2$ $8 \times 3$	16, 24	3	M1 for $40 \div (2 + 3) (= 8)$ oe or $\frac{2}{5}$ oe or $\frac{3}{5}$ oe or for listing at least 3 multiples of 2 and 3 M1 (dep) for "8" $\times$ 2 or "8" $\times$ 3 oe A1 for 16 and 24 in correct places [SC : B2 for 24, 16 if M0 scored] [SC: If M0 scored, B1 may be awarded for just one correct answer, in the correct place]
8	(a)  (b)	15 - 19  Freq polygon through (2, 8), (7, 11), (12, 9), (17, 14) and (22, 18)	1  2	B1 for 15 - 19 oe (eg 15 to 19)  B2 for a complete and correct polygon (ignore any histograms, any lines below a mark of 2 or above a line of 22, but award B1 only if there is a line joining the first to last point)  (B1 for one vertical or one horizontal plotting error OR for incorrect but consistent error in placing the midpoints horizontally (accept end points of intervals) OR for correct plotting of mid-interval values but not joined ) Plotting tolerance $\pm \frac{1}{2}$ square Points to be joined by lines (ruled or hand-drawn but <b>not</b> curves)

Question	Working	Answer	Mark	Notes																
9	$\frac{1}{2} \times 3 \times 4 \times 20$	120	2	M1 for $\frac{1}{2} \times 3 \times 4 \times 20$ A1 cao																
10	$\begin{array}{r} 452 \\ \underline{36} \\ 2712 \end{array}$ $\begin{array}{r} 13560 \\ \underline{16272} \end{array}$  <table border="1" data-bbox="376 774 772 933"> <tr> <td></td> <td><b>400</b></td> <td><b>50</b></td> <td><b>2</b></td> </tr> <tr> <td><b>3</b></td> <td>12000</td> <td>1500</td> <td>60</td> </tr> <tr> <td><b>0</b></td> <td></td> <td></td> <td></td> </tr> <tr> <td><b>6</b></td> <td>2400</td> <td>300</td> <td>12</td> </tr> </table> $12000 + 1500 + 60 + 2400 + 300 + 12 = 16272$		<b>400</b>	<b>50</b>	<b>2</b>	<b>3</b>	12000	1500	60	<b>0</b>				<b>6</b>	2400	300	12	162.72	3	M1 for complete method with relative place value correct. Condone 1 multiplication error, addition not necessary. OR M1 for a complete grid. Condone 1 multiplication error, addition not necessary. OR M1 for sight of a complete partitioning method, condone 1 multiplication error. Final addition not necessary. A2 for 162.72 (A1 (dep on M1) for correct placement of decimal point after final addition (of appropriate values) or for digits 16272 seen) (SC; B1 for attempting to add 36 lots of 4.52)
	<b>400</b>	<b>50</b>	<b>2</b>																	
<b>3</b>	12000	1500	60																	
<b>0</b>																				
<b>6</b>	2400	300	12																	

Question	Working	Answer	Mark	Notes
11	$300 \div 6 = 50$ $300 \div 10 \times 3 = 90$ $300 - 90 - 50$ <p>or</p> $\frac{1}{6} + \frac{3}{10} = \frac{7}{15}$ $\frac{7}{15} \times 300 = 140$ $300 - 140$	160	4	<p>M1 for <math>300 \div 6</math> or 50 seen  M1 for <math>300 \div 10 \times 3</math> oe or <math>30 + 30 + 30</math> or 90 seen  M1 (dep on at least 1 previous M1) for <math>300 - "50" - "90"</math>  A1 cao</p> <p>or</p> <p>M1 for <math>\frac{1}{6} + \frac{3}{10}</math> or <math>\frac{7}{15}</math> oe</p> <p>M1 for "<math>\frac{7}{15}</math>" <math>\times</math> 300 or 140 seen or <math>1 - \frac{7}{15}</math> or <math>\frac{8}{15}</math> oe seen  M1 (dep on at least 1 previous M1) for <math>300 - "140"</math>  or 160 seen or "<math>\frac{8}{15}</math>" <math>\times</math> 300  A1 cao</p>

Question		Working	Answer	Mark	Notes
12		$360 \div 5$ or $180 - (3 \times 180 \div 5)$	72	2	M1 for $360 \div 5$ or $180 - (3 \times 180 \div 5)$ A1 cao
13			2 reasons	2	B2 for 2 out of 3 of these aspects Aspect 1: no time frame, Aspect 2: overlapping, Aspect 3: not exhaustive (B1 for 1 aspect) [SC: B1 for designing a better question identifying at least one aspect]
14	(a)		3, -3, -1	2	B2 for all 3 correct (B1 for 1 or 2 correct)
	(b)		Graph	2	B2 for a fully correct graph or B1 ft for "7 points" plotted correctly $\pm 2$ mm B1 for a smooth curve drawn through their points provided B1 awarded in (a) Note: A straight line drawn from (-1, -3) to (0, -3) gets a maximum of B1
	(c)		-2.3 and 1.3	1	B1 for -2.3 and 1.3 or ft $\pm 2$ mm on a graph with exactly 2 points of intersection with the x-axis.

Question	Working	Answer	Mark	Notes
15	<div style="display: flex; align-items: flex-start;"> <div style="margin-right: 20px;">           2) 180            2) 90            3) 45            3) 15            5) 5            1         </div> <div style="text-align: center;"> <math display="block">  \begin{array}{c}  180 \\  \swarrow \quad \searrow \\  2 \quad 90 \\  \quad \swarrow \quad \searrow \\  \quad 3 \quad 30 \\  \quad \quad \swarrow \quad \searrow \\  \quad \quad 3 \quad 10 \\  \quad \quad \quad \swarrow \quad \searrow \\  \quad \quad \quad 2 \quad 5  \end{array}  </math> </div> </div>	$2 \times 2 \times 3 \times 3 \times 5$	3	<p>M1 for attempt at continual prime factorization (at least two correct divisions); could be shown as a factor tree</p> <p>OR sight of at least one of each 2,3,5 as factors of 180</p> <p>A1 for a fully correct factor tree or 2, 2, 3, 3, 5 which may include 1, but no other numbers</p> <p>A1 for <math>2 \times 2 \times 3 \times 3 \times 5</math> or <math>2^2 \times 3^2 \times 5</math></p> <p>[Note <math>1 \times 2 \times 2 \times 3 \times 3 \times 5</math> or 2,2,3,3,5 or 2.2.3.3.5 do not get the final A1]</p>
16	$\frac{13}{4} \times \frac{8}{3}$	$\frac{26}{3}$	3	<p>M1 for attempt to convert to improper fractions eg. <math>\frac{3 \times 4 + 1}{4}</math> oe or <math>\frac{2 \times 3 + 2}{3}</math> oe seen</p> <p>M1 (dep) for <math>\frac{"13" \times "8"}{4 \times 3}</math> or <math>\frac{104}{12}</math> oe seen</p> <p>A1 for <math>\frac{26}{3}</math> or <math>8\frac{2}{3}</math></p> <p><b>OR</b></p> <p>M1 for <math>3.25 \times 2.66</math> or better</p> <p>M1 for a fully correct multiplication method</p> <p>A1 for 8.66.... (recurring)</p>

Question		Working	Answer	Mark	Notes									
17	(a)		$3(x + 4)$	1	B1 for $3(x + 4)$ Accept $3 \times (x + 4)$ , $(x + 4)3$ and $(x + 4) \times 3$									
	(b)	$8x - 12 = 5x + 7$ $8x - 5x = 7 + 12$ $3x = 19$	$\frac{19}{3}$ oe	3	M1 for $4 \times 2x - 4 \times 3$ or $8x - 12$ seen or for an intent to divide by 4 throughout eg. $\frac{5}{4}x + \frac{7}{4}$ oe seen M1 for a correct method to isolate terms in $x$ and isolate number terms on opposite side of a 4-term equation eg. ' $8x$ ' - $5x = 7 +$ ' $12$ ' or $3x = 19$ seen A1 for $\frac{19}{3}$ oe (accept 6.33 or better)									
	(c)	$y^2 + 5y + 4y + 20$  <table style="border-collapse: collapse; margin-left: 20px;"> <tr> <td style="border-right: 1px solid black; padding: 0 5px;"></td> <td style="padding: 0 5px;"><math>y</math></td> <td style="padding: 0 5px;"><math>+4</math></td> </tr> <tr> <td style="border-top: 1px solid black; border-right: 1px solid black; padding: 0 5px;"><math>y</math></td> <td style="border-top: 1px solid black; padding: 0 5px;"><math>y^2</math></td> <td style="border-top: 1px solid black; padding: 0 5px;"><math>4y</math></td> </tr> <tr> <td style="border-right: 1px solid black; padding: 0 5px;"><math>+5</math></td> <td style="padding: 0 5px;"><math>5y</math></td> <td style="padding: 0 5px;"><math>20</math></td> </tr> </table>		$y$	$+4$	$y$	$y^2$	$4y$	$+5$	$5y$	$20$	$y^2 + 9y + 20$	2	B2 cao (B1 for 4 correct terms with or without signs, or 3 out of no more than 4 terms, with correct signs. The terms may be in an expression or in a table)
	$y$	$+4$												
$y$	$y^2$	$4y$												
$+5$	$5y$	$20$												
	(d)		$4x(2x + 3y)$	2	B2 cao [B1 for $4(2x^2 + 3xy)$ or $x(8x + 12y)$ or $2x(4x + 6y)$ or $2(4x^2 + 6xy)$ or $4x$ (a linear expression in $x$ and $y$ , with just one error); for example $4x(kx + 3y)$ or $4x(2x + ky)$ $k \neq 0$ ]									



Question		Working	Answer	Mark	Notes
18	(a)	$12 \times \frac{6}{4}$	18	2	M1 for sight of $\frac{6}{4}$ oe or $\frac{4}{6}$ oe or $\frac{12}{4}$ oe or $\frac{4}{12}$ oe or a ratio eg. 6:4 oe or decimal eg. 1.5 oe A1 cao
	(b)	$15 \times \frac{4}{6}$	10	2	M1 for $15 \times \frac{4}{6}$ oe or $\frac{15}{6} \times 12$ oe A1 cao
19		$4000 - \frac{10}{100} \times 4000 = 3600$ $3600 - \frac{10}{100} \times 3600$	3240	3	M1 for $4000 - \frac{10}{100} \times 4000$ or $0.9 \times 4000$ oe or 3600 or 400 or 3200 or 800 seen M1 (dep) "3600" - $\frac{10}{100} \times$ "3600" or "3600" $\times 0.9$ oe A1 cao or M2 for $4000 \times 0.9^2$ (M1 for $4000 \times 0.9^3$ ) A1 cao  [SC: B2 for an answer of £4840, with or without working]

Question		Working	Answer	Mark	Notes
20	(a)		$a^2(c+b)$ $4abc$	2	B1 for $a^2(c+b)$ B1 for $4abc$ [-1 for each additional incorrect answer, up to a minimum of 0]
	(b)	$8 \times 100 \times 100 \times 100$	8 000 000 or $8 \times 10^6$ or 8 million	2	M1 for sight of $10^6$ oe or $100 \times 100 \times 100$ or $200 \times 200 \times 200$ A1 for 8 000 000 or $8 \times 10^6$
21		$6x + 4y = 16$ $6x + 15y = -6$ $-11y = 22$ $6x + 4 \times -2 = 16$  <b>Alternative method</b> $x = \frac{8-2y}{3}$ $2\left(\frac{8-2y}{3}\right) + 5y = -2$ $16 - 4y + 15y = -6$ $11y = -22$  $x = \frac{8-2 \times -2}{3}$	$x = 4, y = -2$	4	M1 for correct process to eliminate either x or y (condone one arithmetic error) A1 for either $x = 4$ or $y = -2$  M1 (dep on 1 <sup>st</sup> M1) for correct substitution of their found variable OR M1 (indep of 1 <sup>st</sup> M1 for a correct process to eliminate the other variable (condone one arithmetic error) A1 cao for both $x = 4$ and $y = -2$  [SC: B1 for $x=4$ or $y=-2$ if M0 scored]

Question		Working	Answer	Mark	Notes
22	(a)		$20 < n \leq 30$	1	B1 for $20 < n \leq 30$ Accept 20 to 30, 20 - 30 oe but <b>not</b> 26 Accept an indication of chosen interval on the diagram (circling) if no answer on the answer line
	(b)		16,42,65,75,80	1	B1 cao
	(c)		Points plotted and joined	2	B1 ft for at least 4 of "5 points" plotted correctly $\pm 2$ mm at end of interval dep on sensible table (condone 1 addition error) B1 ft (dep on previous B1) for points joined by curve or line segments provided no gradient is negative - ignore any part of graph outside range of their points (SC B1 if 4 or 5 pts plotted not at end but consistent within each interval <b>and</b> joined)
	(d)(i)		28 - 30	3	B1 for an answer in the range 28 - 30 or from "cf graph"
	(ii)		15 - 17		M1 for horizontal lines drawn at cf = 20 and cf = 60 oe and vertical lines drawn to 'x'-axis or 'correct' marks drawn on 'x'-axis only or for UQ = 36 - 38 and LQ = 20.5 - 23 or ft "cf graph" A1 For answer in the range of 15-17 or ft from "cf graph"

Question		Working	Answer	Mark	Notes
23		Gradient = $\frac{10 - -2}{3 - 0}$	$y = 4x - 2$	3	M1 for gradient = $\frac{10 - -2}{3 - 0}$ oe or $(y =) 4x + c$ or a right angle triangle with sides 12 and 3 shown M1 for $(y =) mx - 2$ , $m \neq 0$ or $10 = 3m + c$ or $-2 = c$ (but not 'y-intercept = -2')  A1 for $y = 4x - 2$ oe [the y must be included in the equation]
24	(i)		1	1	B1 cao
	(ii)		8	1	B1 for 8 or -8 or $\pm 8$
	(iii)		$\frac{4}{9}$	2	M1 for $\left(\frac{8}{27}\right)^{\frac{2}{3}}$ oe or $\left(\frac{3}{2}\right)^{-2}$ oe or $\left(\frac{2}{3}\right)^2$ oe or $\left(\frac{1}{\sqrt[3]{27/8}}\right)^2$ or better or $\frac{9}{4}$ oe seen A1 cao

Question	Working	Answer	Mark	Notes
25	<p>(a) <math>x^2 + (x+2)^2 = (x+4)^2</math></p> $x^2 + x^2 + 4x + 4 = x^2 + 8x + 16$ $x^2 + 4x - 8x + 4 - 16 = 0$ <p>b)(i) <math>(x-6)(x+2) = 0</math>  <math>x-6=0</math> or <math>x+2=0</math> or</p> $x = \frac{- -4 \pm \sqrt{(-4)^2 - 4 \times 1 \times -12}}{2 \times 1}$ $x = \frac{4 \pm \sqrt{64}}{2}$ <p><math>(x-2)^2 - 16 = 0</math>  <math>x-2 = \pm\sqrt{16}</math>  <math>x = 2 \pm\sqrt{16}</math></p> <p>(ii)</p>	<p>Proof</p> <p>6, -2</p> <p>10</p>	<p>3</p> <p>4</p>	<p>M1 for <math>x^2 + (x+2)^2 = (x+4)^2</math> oe  A1 for <math>x^2+x^2+4x+4 = x^2+8x+16</math>  A1 for completing the proof</p> <p>M1 for <math>(x \pm 6)(x \pm 2) (= 0)</math></p> <p>OR <math>\frac{- -4 \pm \sqrt{(-4)^2 - 4 \times 1 \times -12}}{2 \times 1}</math>  allow <math>\pm 4</math> for b and <math>\pm 12</math> for c</p> <p>OR <math>(x-2)^2 - 16 = 0</math></p> <p>A1 <math>x = 6</math> cao  A1 <math>x = -2</math> cao  [SC: B1 for 1 correct solution if MO scored]  B1 ft from (b)(i), provided x is a positive value  [Note: an answer of 10 and 2 gets B0]</p>

Question	Working	Answer	Mark	Notes
26	$\left(\frac{3}{10} \times \frac{2}{9}\right) + \left(\frac{2}{10} \times \frac{1}{9}\right) + \left(\frac{5}{10} \times \frac{4}{9}\right) = \frac{6+2+20}{90}$	$\frac{28}{90} \text{ oe}$	4	<p>B1 for <math>\frac{2}{9}</math> (orange) or <math>\frac{1}{9}</math> (red) or <math>\frac{4}{9}</math> (yellow) seen as 2<sup>nd</sup> probability</p> <p>M1 for <math>\left(\frac{3}{10} \times \frac{2}{9}\right)</math> or <math>\left(\frac{2}{10} \times \frac{1}{9}\right)</math> or <math>\left(\frac{5}{10} \times \frac{4}{9}\right)</math></p> <p>M1 for <math>\left(\frac{3}{10} \times \frac{2}{9}\right) + \left(\frac{2}{10} \times \frac{1}{9}\right) + \left(\frac{5}{10} \times \frac{4}{9}\right)</math></p> <p>A1 for <math>\frac{28}{90}</math> oe</p> <p><b>Alternative scheme for replacement</b></p> <p>M1 for <math>\left(\frac{3}{10} \times \frac{3}{10}\right)</math> or <math>\left(\frac{2}{10} \times \frac{2}{10}\right)</math> or <math>\left(\frac{5}{10} \times \frac{5}{10}\right)</math></p> <p>M1 for <math>\left(\frac{3}{10} \times \frac{3}{10}\right) + \left(\frac{2}{10} \times \frac{2}{10}\right) + \left(\frac{5}{10} \times \frac{5}{10}\right)</math></p> <p>No further marks may be awarded</p>

Question	Working	Answer	Mark	Notes
27	<p> <math>PQT = 58^\circ</math> (Alternate segment theorem)  <math>QTP = (180 - 58)/2 (= 61^\circ)</math> (Isosceles triangle)  <math>OTQ = 61 - (90 - 58)</math> (Angle between tangent and radius) </p> <p>           Alternative:  <math>OTP = 90 - 58 (= 32^\circ)</math> (angle between radius and tangent)  <math>OPT = OPT = 32^\circ</math> (isosceles triangle)  <math>POT = 180 - 32 - 32 = 116^\circ</math> (angles in a triangle)  <math>PQT = 116 \div 2 = 58^\circ</math> (angle at centre = <math>2 \times</math> angle at circumference)  <math>QTP = QPT = (180 - 58)/2 (= 61^\circ)</math> (Isosceles triangle) </p> <p> <math>OTQ = 61 - 32</math> </p>	29	5	<p>           M1 for <math>PQT = 58^\circ</math> or <math>\frac{1}{2} (180 - 2(90 - 58))</math>            M1 (dep) for <math>QPT</math> or <math>QTP = (180 - "58")/2</math>            A1 for <math>OTQ = 29^\circ</math> </p> <p>           B2 for fully correct reasons            (B1 for alternate segment theorem or equivalent circle theorems leading to <math>PQT</math>) </p>